



While deep vein thrombosis (DVT) and PE are relatively common clinically, they are rare in the absence of elevated blood D-Dimer levels and certain specific risk factors. Imaging, particularly computed tomography (CT) pulmonary angiography, is a rapid, accurate, and widely available test, but has limited value in patients who are very unlikely, based on serum and clinical criteria, to have significant value. Imaging is helpful to confirm or exclude PE only for such patients, not for patients with low pre-test probability of PE. Source: American College of Radiology (ACR). Wording reflects that of the Radiology recommendation, other societies have similar recommendations, some explicitly recommended D-Dimer testing prior to imaging.



There is no indication to order these tests, and there is no benefit to be derived in obtaining them in someone that does not have any history of bleeding or abnormal clotting and in the absence of any family history. This testing is not a part of the infertility workup. Furthermore, the testing is costly, and there are risks associated with the proposed treatments, which would also not be indicated in this routine population. *Source:* 

Torbicki A, Perrier A, Konstantinides S, et al. Guidelines on the diagnosis and management of acute pulmonary embolism: the Task Force for the Diagnosis and Management of Acute Pulmonary Embolism of the European Society of Cardiology (ESC). Eur Heart J. 2008;29(18):2276–315.

Neff MJ. ACEP releases clinical policy on evaluation and management of pulmonary embolism. American Family Physician 2003;68(4):759-60.

Stein PD, Woodard PK, Weg JG, Wakefield TW, Tapson VF, Sostman HD, Sos TA, Quinn DA, Leeper KV, Hull RD, Hales CA, Gottschalk A, Goodman LR, Fowler SE, Buckley JD. Diagnostic pathways in acute pulmonary embolism: recommendations of the PIOPED II Investigators. Radiology 2007;242(1):15–21.

Lockwood C, Wendel G; Committee on Practice Bulletins-Obstetrics. Practice bulletin no. 124: inherited thrombophilias in pregnancy. Obstet Gynecol. 2011 Sept;118(3):730-40.

Casadei L, Puca F, Privitera L, Zamaro V, Emidi E. Inherited thrombophilia in infertile women: implication in unexplained infertility. Fertil Steril. 2010 Jul;94(2):755-7.

The Practice Committee of the American Society for Reproductive Medicine. Diagnostic evaluation of the infertile female: a committee opinion. Fertil Steril. 2012 Aug;98:302-7.

Baglin T, Gray E, Greaves M, Hunt B, Keeling D, Machin S, Mackie I, Makris M, Nokes T, Perry D, Talt RC, Walker I, Watson H. Clinical guidelines for testing for heritable thrombophilia. Br J Haematol. 2010;149:209–20.

Salisbury AC, Reid KJ, Alexander KP, Masoudi FA, Lai SM, Chan PS, Bach RG, Wang TY, Spertus JA, Kosiborod M. Diagnostic blood loss from phlebotomy and hospital-acquired anemia during Acute Myocardial Infarction. Arch Intern Med [Internet]. 2011 Oct 10 [cited 2012 Sep 4]:171(18):1646-1653.

Thavendiranathan P, Bagai A, Ebidia A, Detsky AS, Choudhry NK. Do blood tests cause anemia in hospitalized patients?: The effect of diagnostic phlebotomy on hemoglobin and hematocrit levels. J Gen Intern Med [Internet]. 2005 June [cited 2012 Sep 4]:20(6):520–524.

Stuebing EA, Miner TJ. Surgical vampires and rising health care expenditure: reducing the cost of daily phlebotomy. Arch Surg [Internet]. 2011 May [cited 2012 Sep 4];146(5):524-7.

AABB. Guidelines for patient blood management and blood utilization. Bethesda (MD): AABB; 2011. 52 p.

Lin DM, Lin ES, Tran MH. Efficacy and safety of erythropoietin and intravenous iron in perioperative blood management: a systematic review. Transfus Med Rev. 2013 Oct;27(4):221-34.

Friedman AJ, Chen Z, Ford P, Johnson CA, Lopez AM, Shander A, Waters JH, van Wyck D. Iron deficiency anemia in women across the life span. J Womens Health (Larchmt). 2012 Dec;21(12):1282-9.

Phurrough S, Cano C, Dei Cas R, Ballantine L, Carino T; Centers for Medicare and Medicaid Services. Decision memo for positron emission tomography (FDG) for solid tumors (CAG-00181R4). Baltimore